SuperNova / Acceleration Probe
DOE Review:
Science Overview

Lawrence Berkeley Laboratory
July 2002

Saul Perlmutter

0. Background: cosmology/dark energy measurements from supernovae.
1. A really hard goal: $w'(z)$.
2. An exhaustive list of systematics.
3. An unusual tool for cosmology.
4. How this tool can address systematics.
5. Project status and reviews.


FAINTER
(Farther)
(Further back in time)
Type Ia Supernovae
Supernova Cosmology Project
Calan/Tololo Supernova Survey
Open
Flat
Closed
magnitude
redshift
MORE REDSHIFT
(More total expansion of universe since light left the Standard Candle)


FAINTER
(Farther)
(Further back in time)
Type Ia Supernovae
Λ -- cosmological constant
Open
Flat
Closed
Supernova Cosmology Project
Calan/Tololo Supernova Survey
Accelerating Universe
Decelerating Universe
magnitude
redshift
MORE REDSHIFT
(More total expansion of universe since light left the Standard Candle)



Perlmutter *et al.*
(Ap.J. 1999)
astro-ph/98 12133


Perlmutter, et al. (1999)
Jaffe et al. (2000)
Bahcall et al. (2000)
No Big Bang
Supernovae
CMB
Clusters
expands forever
recollapses eventually
closed
flat
open
vacuum energy density
(cosmological constant)
mass density
3
2
1
0
-1
0
1
2
3

What's wrong with a non-zero vacuum energy / cosmological constant?

Two coincidences:

- ***Why so small?***

  Might expect $\frac{\Lambda}{8\pi G} \sim m^4_{\text{Planck}}$

  This is off by ~120 orders of magnitude!

- ***"Why now?"***

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3}(\rho + 3p)$$

MATTER: $p = 0 \rightarrow \rho \propto R^{-3}$

VACUUM ENERGY: $p = -\rho \rightarrow \rho \propto constant$

What's wrong with a non-zero vacuum energy / cosmological constant?

Two coincidences:

- ***Why so small?***

  Might expect $\frac{\Lambda}{8\pi G} \sim m^4_{\text{Planck}}$

  This is off by ~120 orders of magnitude!

- ***"Why now?"***

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3}(\rho + 3p)$$

MATTER: $p = 0 \rightarrow \rho \propto R^{-3}$

VACUUM ENERGY: $p = -\rho \rightarrow \rho \propto constant$



What's wrong with a non-zero vacuum energy / cosmological constant?

Two coincidences:

- ***Why so small?***

  Might expect $\frac{\Lambda}{8\pi G} \sim m^4_{\text{Planck}}$

  This is off by ~120 orders of magnitude!

- ***"Why now?"***

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3}(\rho + 3p)$$

MATTER: $p = 0 \rightarrow \rho \propto R^{-3}$

VACUUM ENERGY: $p = -\rho \rightarrow \rho \propto constant$

What are the alternatives?

New Physics: "Dark energy":

Dynamical scalar fields, "quintessence",...

General Equation of State:

$$p = w\rho \rightarrow \rho \propto R^{-3(1+w)}$$

and w can vary with time

Unknown Component, Ω_u, of Energy Density



1. A Really Hard Goal: $w'(z)$

A measurement of the expansion history of the universe with enough accuracy that a measurement of a *change* in the properties of the dark energy, e.g. $w' \neq 0$, would be trusted.


FAINTER
(Farther)
(Further back in time)
Type Ia Supernovae
Λ -- cosmological constant
Open
Flat
Closed
Supernova Cosmology Project
Calan/Tololo Supernova Survey
Accelerating Universe
Decelerating Universe
magnitude
redshift
MORE REDSHIFT
(More total expansion of universe since light left the Standard Candle)

1998: Acceleration



New HST data


Supernova Cosmology Project
Knop, et al. *Preliminary*
fainter
Δmag
1.0
0.5
0.0
-0.5
-1.0
0.0
0.5
1.0
1.5
2.0
redshift
Empty
Accelerating (Λ)
Open (no Λ)
Flat (no Λ)

New HST data


Supernova Cosmology Project
Knop, et al. *Preliminary*
fainter
Δmag
1.0
0.5
0.0
-0.5
-1.0
0.0
0.5
1.0
1.5
2.0
redshift
Example (extreme?)
Dust or Evolution
Empty
Accelerating (Λ)
Open (no Λ)
Flat (no Λ)

Supernovae probing the *deceleration* era with IR measurements




Simulated
Future HST Data
Example (extreme?)
Dust or Evolution
Empty
Accelerating (Λ)
Open (no Λ)
Flat (no Λ)
SN 1997ap
SN 1998eq
SN 1997ff
Δmag
fainter
1.0
0.5
0.0
-0.5
-1.0
0.0
0.5
1.0
1.5
2.0
redshift


Simulated
Future HST Data
fainter
Δmag
1.0
0.5
0.0
-0.5
-1.0
0.0
0.5
1.0
1.5
2.0
redshift
Example (extreme?)
Dust or Evolution
Empty
Accelerating (Λ)
Open (no Λ)
Flat (no Λ)
SN 1997ap
SN 1998eq
SN 1997ff

Current statistical uncertainties from supernova measurements are almost good enough that they are limited by systematic uncertainties:

Uncertainties on Ω_M or Ω_Λ in flat cosmology:

Statistical	
high-redshift SNe	0.05
low-redshift SNe	0.065
Total	***0.085***
Systematic Total	
identified entities/processes	***0.05***

Significant advances in these measurement uncertainties will require **much better constraints on the systematics.**

Perlmutter et al. (1999)

2. An exhaustive list of systematics.

What makes the supernova measurement special?

An exhaustive accounting of sources of SN systematic uncertainties:

SN Ia Evolution

- shifting distribution of progenitor mass/metallicity/C-O
- shifting distibution of SN physics params:
  - -- amount of Nickel fused in explosion
  - -- distribution of Nickel
  - -- kinetic energy of explosion
  - -- opacity of atmosphere's inner layers
  - -- metallicity

Gravitational Lensing (de)amplification

Dust/Extinction

- dust that reddens
- evolving gray dust
  - -- clumpy
  - -- homogeneous
- Galactic extinction model

Observational biases

- Malmquist bias differences
- non-SN Ia contamination
- K-correction uncertainty
- color zero-point calibration

Perlmutter et al. (1999)

Supernova Demographics



Gravitational Lensing in a Clumpy Universe

"Weak Lensing" Approx:
Power spectrum of mass density in relatively smooth universe.





"Cross-Filter" K corrections

Kim, Goobar, & Perlmutter (1995)



3. SN Ia: An unusual tool for cosmology.





Kim, *et al.* (1997)

The time series of spectra is a “CAT Scan” of the Supernova

-14 days





maximum





+10 days





+20 days





Time Series of Low-Redshift and High-Redshift Spectra



Control of Evolution Systematics:
Matching Supernovae



SNAP SuperNova Acceleration Probe



SNAP SuperNova Acceleration Probe



Control of Evolution Systematics:
Matching Supernovae



Host galaxy morphology, from space-based imaging with HST / STIS

Result from elliptical host galaxy subsample agrees with flat, $\Omega_\Lambda = 0.72$ result from whole dataset.
(Elliptical best flat universe fit: $\Omega_\Lambda = 0.58 \pm 0.2$)







galaxy type	dispersion from flat, $\Omega_\Lambda = 0.72$
● Elliptical: E/S0	$\sigma = 0.19$ mag
△ Spiral: Sa/Sb/Sc	$\sigma = 0.27$ mag
✳ Late/Irregular: Scd/Irr	$\sigma = 0.30$ mag



Ellis et al. (astro-ph/0011369)
Supernova Cosmology Project

4. How this tool can address systematics.

SNAP SuperNova Acceleration Probe



- **~2 m aperture telescope**
  Can reach very distant SNe.

- **1 square degree mosaic camera, ~1 billion pixels**
  Efficiently studies large numbers of SNe.

- **0.35um -- 1.7um spectrograph**
  Detailed analysis of each SN.

SNAP: observing supernovae with lightcurves & spectra

$z = 0.8$



$z = 1.0$





$z = 1.2$





$z = 1.4$





$z = 1.6$





An exhaustive accounting of sources of SN systematic uncertainties:

SN Ia Evolution

- shifting distribution of
  progenitor mass/metallicity/C-O
- shifting distibution of SN physics params:
  -- amount of Nickel fused in explosion
  -- distribution of Nickel
  -- kinetic energy of explosion
  -- opacity of atmosphere's inner layers
  -- metallicity

Gravitational Lensing (de)amplification

Dust/Extinction

- dust that reddens
- evolving gray dust
  -- clumpy
  -- homogeneous
- Galactic extinction model

Observational biases

- Malmquist bias differences
- non-SN Ia contamination
- K-correction uncertainty
- color zero-point calibration

Perlmutter et al. (1999)

Data Sheets
for each SN

SN
0001

Host galaxy morphology



Neighboring galaxy distribution



Lightcurve



Spectrum

SN
0002



Lightcurve



Spectrum

SN
2541

bution



Lightcurve



Spectrum

1

Sort into Like Subsets

Group A:
* Si II in spectrum: type Ia
* elliptical host
* bright UV: low metallicity
* fast rise time: low Ni56 mass
* spectral feature velocities
  9000 < v < 10000 km/s

Group B:
* Si II in spectrum: type Ia
* in core of late-type spiral host
* faint UV: high metallicity
* fast rise time: low Ni56 mass
* spectral feature velocities
  9000 < v < 10000 km/s

Group C:
* Si II in spectrum: type Ia
* in outskirts of late-type spiral host
* bright UV: low metallicity
* long rise time: high Ni56 mass
* spectral feature velocities
  8000 < v < 9500 km/s

Gro
* Si II i
* in cor
* bright
* short
* spectr
  8000



2

Each subset gets its own extinction-corrected Hubble diagram:

Group A:
* Si II in spectrum: type Ia
* elliptical host
* bright UV: low metallicity
* fast rise time: low Ni56 mass
* spectral feature velocities
  9000 < v < 10000 km/s

Group B:
* Si II in spectrum: type Ia
* in core of late-type spiral host
* faint UV: high metallicity
* fast rise time: low Ni56 mass
* spectral feature velocities
  9000 < v < 10000 km/s

Group C:
* Si II in spectrum: type Ia
* in outskirts of late-type spiral host
* bright UV: low metallicity
* long rise time: high Ni56 mass
* spectral feature velocities
  8000 < v < 9500 km/s

• • •

Gro
* Si II i
* in cor
* bright
* short
* spectr
  8000



2 Each subset gets its own extinction-corrected Hubble diagram:



Combine into one Hubble diagram

with magnitude difference from $z = 0.5$



3

Break Hubble diagram into z slices to study lensing (de)amplification distribution:



3 Fit/average lensing distributions to construct redshift-binned Hubble diagram:



Example Measurement Requirements for Each Step

Sort into Like Subsets

1

Spectrum:

Si II feature $\lambda/\Delta\lambda \sim 100$ resolution

UV features 5σ per bin

Lightcurve:

Rise time 3σ measurement
3.8 mag before max

Peak fit 15σ measurement
2 mag after max

Image:

Host galaxy morphology <0.1" dithered resolution

Extinction-corrected Hubble diagram

2

Spectrum & Lightcurve:

Cross-wavelength calibrated colors for photometry and spectroscopy from near-UV to near-IR (0.35 -- 1.7 μm)

Correct for lensing distributions

3

Image quality:

<0.1" dithered resolution for neigboring galaxy gravitational lensing map

Redshift range & statistics:
>~50 SNe per bin to obtain lensing distribution


SNAP
LSST
with NIR camera added
restframe B
restframe V
z=0.8
Band-5
Band-6
z=1.0
Band-5
Band-7
z=1.2
Band-6
Band-7
z=1.4
Band-6
Band-8
z=1.6
flux
-50
0
50
100
150
date (days)

Ground:LSST/VLT
9 hours
OH Suppression

Space: SNAP

Multi-object: No AO

Single-object: With AO

$z = 1.0$ *Water*

Si II

Si II

$z = 1.2$

Si II

$z = 1.4$

Si II

$z = 1.6$

Si II



Ground:LSST/VLT
9 hours
OH Suppression

Space: SNAP

Multi-object: No AO

Single-object: With AO

$z = 1.0$ *Water*

Si II

Si II

$z = 1.2$

Si II

$z = 1.4$

Si II

$z = 1.6$

Si II



SNAP SuperNova Acceleration Probe

Current ground-based data
compared with binned simulated SNAP data
and a sample of Dark Energy models.



based on
Weller & Albrecht (2001)

SNAP SuperNova Acceleration Probe

Binned simulated SNAP data compared with Dark Energy models currently in the literature.



based on

Weller & Albrecht (2001)

SNAP SuperNova Acceleration Probe

Binned simulated SNAP data
compared with Dark Energy models.



based on
Weller & Albrecht (2000)

How the uncertainty improves as we extend the redshift range.



SNAP SuperNova Acceleration Probe

Primary Science Mission

Requiring complementary measurements of cosmological parameters, Dark Matter, Dark Energy,...

Type Ia supernova calibrated candle:
Hubble diagram to $z = 1.7$

Type II supernova expanding photosphere:
Hubble diagram to $z = 1$ and beyond.

Weak lensing:
Direct measurements of $P(k)$ vs z
Mass selected cluster survey vs z

Strong lensing statistics: Ω_Λ
10x gains over ground based optical resolution, IR channels + depth.

Galaxy clustering:
$W(\Theta)$ angular correlation vs redshift from 0.5 to 3.0


Perlmutter, et al. (1999)
Jaffe et al. (2000)
Bahcall et al. (2000)
No Big Bang
Supernovae
CMB
Clusters
expands forever
recollapses eventually
closed
flat
open
vacuum energy density
(cosmological constant)
mass density
3
2
1
0
-1
0
1
2
3


Perlmutter, et al. (1999)
Jaffe et al. (2000)
Bahcall et al. (2000)
No Big Bang
Supernovae
SNAP
Target Uncertainty
CMB
Clusters
expands forever
recollapses eventually
closed
flat
open
vacuum energy density
(cosmological constant)
mass density
3
2
1
0
-1
0
1
2
3


Perlmutter, et al. (1999)
Jaffe et al. (2000)
Bahcall et al. (2000)
No Big Bang
Supernovae
It is possible that SNe will give
a result that disagrees
with the flat universe
measurement from the CMB.
CMB
Clusters
expands forever
recollapses eventually
closed
flat
open
vacuum energy density
(cosmological constant)
mass density
3
2
1
0
-1
0
1
2
3

SNAP SuperNova Acceleration Probe

5. Project status & reviews.

SNAP SuperNova Acceleration Probe

Active members of the SNAP collaboration

G. Aldering, C. Bebek, J. Bercovitz, W. Carithers, C. Day, S. Deustua*, R. DiGennaro, D. Groom, S. Holland, D. Huterer*, A. Karcher, A. Kim, W. Kolbe, B. Krieger, R. Lafever, M. Levi, E. Linder, S. Loken, P. Nugent, H. Oluseyi, S. Perlmutter, K. Robinson, A. Spadafora, J. Walder (Lawrence Berkeley National Laboratory)

M. Bester, E. Commins, G. Goldhaber, S. Harris, P. Harvey, H. Heetderks, M. Lampton, J. Lamoureux, D. Pankow, C. Pennypacker, R. Pratt, M. Sholl, G. F. Smoot (UC Berkeley)

C. Akerlof, G. Bernstein*, D. Levin, T. McKay, S. McKee, M. Schubnell, G. Tarle , A. Tomasch (U. Michigan)

R. Ellis, R. Massey*, J. Rhodes, A. Refregier* (CalTech)

C. Bower, N. Mostek, J. Musser, S. Mufson (Indiana)

A. Fruchter (STScI)

P. Astier, E. Barrelet, A. Bonnissent, A. Ealet, J-F. Genat, R. Malina, R.Pain, E. Prieto, G. Smadja, D. Vincent (France: CNRS/IN2P3, CNRS/INSU)

R. Amanullah, L. Bergström, M. Eriksson, A. Goobar, E. Mörtsell (U. Stockholm)

Samuel Silver Space Sciences Laboratory



IN2P3

** Affiliated with the listed institution for organizational purposes*

SNAP SuperNova Acceleration Probe

Project Chronology: Reviews & Studies

First public presentation of idea at Fermilab "Inner Space/Outer Space" symposium.	**May 1999**
SAGENAP review 1 for DOE & NSF-Physics	**March 2000**
NASA Structure and Evolution of the Universe (SEU) subcomittee	**Nov 2000**
NAS/NRC Committee on Astronomy and Astrophysics	**Dec 2000**
DOE-HEP R&D review	**Jan 2001**
DOE HEPAP	**March 2001**
NASA Integrated Mission Design Center	**June 2001**
NAS/NRC Committee on the Physics of the Universe	**July 2001**
CNES (France Space Agency)	**Nov 2001**
NASA/SEU Strategic Planning Panel	**Dec 2001**
NASA Instrument Synthesis & Analysis Lab	**Dec 2001**
SAGENAP review 2	**March 2002**
CNES review (scheduled)	**Oct 2002**

SNAP SuperNova Acceleration Probe

Project History and Status

Peer review of science (260 page proposal) by
DOE & NSF's SAGENAP panel, March 2000:

Strong endorsement of science,
and recommendation for study funding.

DOE/Science & R&D Review (Jan 2001):

"SNAP is a science-driven project with compelling scientific goals."
"SNAP will have a unique ability to measure the variation in the equation of state of the universe."
"Implications for particle physics: We believe that it is not an overstatement to say that the Type Ia supernova measurements will uniquely address issues at the very heart of the field…"

SNAP SuperNova Acceleration Probe

Three National Prioritization Panels' Recommendations

Astronomy & Astrophysics

NRC Decadal Survey:

"One of the most exciting developments of the past decade has been the discovery that the cosmological constant may not be zero — our universe appears to be filled with dark energy."

"The committee identified several key problems that are particularly ripe for advances in the coming decade. These problems are … properties of the universe: the amount and distribution of its matter and energy, its age, and the history of its expansion."

SNAP was formulated after the Decadal Survey's data collection phase.

Physics

HEPAP 20-Year Planning Report:

"Modern cosmology is closely connected with particle physics. For example, cosmological measurements of dark energy and particle dark matter have direct implications for particle physics."

"Dark energy can be probed by a number of techniques. Among the most powerful are measurements of the expansion rate of the universe from observations of Type Ia supernovae."

Report gives a strong endorsement for continued development of SNAP.

Intersection of Physics and Astronomy

NRC Committee on the Physics of the Universe:

"Deciphering the nature of dark matter and dark energy is one of the most important goals in the physics of the universe. Resolving both puzzles is key to advancing our understanding not only of cosmology but also fundamental physics."

"Observations of distant supernovae can probe the detailed expansion history directly back to redshifts of around 2.... Large-field-of-view telescopes are needed to find larger and more uniform samples of supernovae."

Committee reviewed SNAP in July 2001 as part of their Phase II study of specific projects (released this spring).

Report of the NRC's Commitee on the Physics of the Universe (The Turner Panel)

Recommendation:

"The discovery that the expansion of the universe is speeding up and not slowing down has revealed the presence of a mysterious new energy form that accounts for two-thirds of all the matter and energy in the universe. Because of its diffuse nature, it can only be probed through its effect on the expansion of the universe.

"The NRC's most recent astronomy decadal survey has recommended building the Large Synoptic Survey Telescope to study transient phenomena in the universe; it will also have significant ability to probe dark energy. To fully characterize the expansion history and probe the dark energy will require a wide-field telescope in space (such as the Supernova/Acceleration Probe)."

Recommendation: "The Committee further recommends that NASA and DOE work together to construct a wide-field telescope in space to determine the expansion history of the universe and fully probe the nature of the dark energy."

SNAP SuperNova Acceleration Probe

Science Goals for
The First Wide-field Survey in Space

A Resource for the Science Community:
The *only* wide-field deep survey in space -- with HST resolution.

- SNAP main survey will be 4000x larger (and somewhat deeper) than the biggest HST deep survey, the ACS survey
- Complementary to NGST: target selection for rare objects
- Can survey 3000 sq. deg in a year to I = 29 or J = 28 (AB mag) .



Archive data distributed

Guest Survey Program

Whole sky can be observed every few months

SNAP SuperNova Acceleration Probe

Grass-roots Support

January 2001 AAS meeting
1 dedicated session, 9 talks, 4 posters
January 2002 AAS meeting
2 dedicated sessions, 18 talks, 7 posters:

Oral Session 111. Science with Wide Field Imaging in Space:

The Astronomical Potential of Wide-field Imaging from Space	S. Beckwith (Space Telescope Science Institute)
Galaxy Evolution: HST ACS Surveys and Beyond to SNAP	G. Illingworth (UCO/Lick, U. of California)
Studying Active Galactic Nuclei with SNAP	P.S. Osmer (OSU), P.B. Hall (Princeton/Catolica)
Distant Galaxies with Wide-Field Imagers	K. M. Lanzetta (State U. of NY at Stony Brook)
Angular Clustering and the Role of Photometric Redshifts	A. Conti, A. Connolly (University of Pittsburgh)
SNAP and Galactic Structure	I. N. Reid (STScI)
Star Formation and Starburst Galaxies in the Infrared	D. Calzetti (STScI)
Wide Field Imagers in Space and the Cluster Forbidden Zone	M. E. Donahue (STScI)
An Outer Solar System Survey Using SNAP	H.F. Levison, J.W. Parker (SwRI), B.G. Marsden (CfA)

Oral Session 116. Cosmology with SNAP:

Dark Energy or Worse	S. Carroll (University of Chicago)
The Primary Science Mission of SNAP	S. Perlmutter (Lawrence Berkeley Laboratory),
The Supernova Acceleration Probe:mission design & core survey	T. A. McKay (University of Michigan)
Sensitivities for Future Space- and Ground-based Surveys	G. M. Bernstein (Univ. of Michigan)
Constraining the Properties of Dark Energy using SNAP	D. Huterer (Case Western Reserve University)
Type Ia Supernovae as Distance Indicators for Cosmology	D. Branch (U. of Oklahoma)
Weak Gravitational Lensing with SNAP	A. Refregier (Cambridge), Richard Ellis (Caltech)
Strong Gravitational Lensing with SNAP	R. D. Blandford, L. V. E. Koopmans, (Caltech)
Strong lensing of supernovae	D.E. Holz (ITP, UCSB)

Poster Session 64. Overview of The Supernova/Acceleration Probe:

Supernova / Acceleration Probe: An Overview	M. Levi (LBNL)
The SNAP Telescope	M. Lampton (UCB)
SNAP: Science with Wide Deep Fields in Space	E. Linder (LBNL)
SNAP: An Integral Field Spectrograph for Supernova Identification	R. Malina (Marseille,INSU), A. Ealet
Supernova / Acceleration Probe: GigaCAM - A Billion Pixel Imager	C. Bebek (LBNL)
Supernova / Acceleration Probe: Cosmology with Type Ia Supernovae	A. Kim (LBNL)
Supernova / Acceleration Probe: Education and Outreach	S. Deustua (LBNL)

SNAP SuperNova Acceleration Probe

Binned simulated SNAP data compared with Dark Energy models currently in the literature.



based on

Weller & Albrecht (2001)

CMB data before BOOMERANG and MAXIMA





Expansion History of the Universe



END